

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any problems that occur due to the contents of the multilingual product information.

You should read this written explanation before use.
Please keep it so that you can read it when you need to.



大幸薬品

セイロガン糖衣A®

2nd Class OTC drug

Gastrointestinal
medicine

Seirogan Toi-A

The gastrointestinal medicine **Seirogan Toi-A** (sugar-coated tablet), is part of a product lineup that includes Seirogan, a household medicine that has been used for over a century. Japanese Pharmacopeia (JP) wood creosote, the main active ingredient of this medicine, regulates the water balance in the intestines without stopping bowel peristalsis, and returns the intestinal environment to its normal condition. It is especially effective for soft stool and diarrhea caused by food poisoning, water poisoning and indigestion, and for soft stool and diarrhea due to stress and cold. This medicine is an odorless, easy-to-swallow white tablet, containing crude drugs such as JP wood creosote, JP powdered geranium herb, and dry phellodendron bark extract.

[Precautions]

When not to use the product



[If you do not follow these instructions, the current symptoms may worsen, or adverse reactions/incidents may occur.]

1. This product should not be taken by the following persons.

- (1) Patients who have had an allergic symptom (rash, redness, itching, oedema, etc.) to this drug or its ingredients
- (2) Patients undergoing dialysis

2. Do not take this medicine for a long time.

Consultation



1. The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.

- (1) Patients undergoing medical treatment from a physician
- (2) Those who have diarrhea with a fever, bloody stool or persistent mucous stool
- (3) Pregnant women, women suspected of being pregnant, or nursing women
- (4) The elderly
- (5) Patients who have experienced allergic symptoms associated with drugs or cosmetics, etc.
- (6) Patients with liver disease and/or renal disease

2. If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this package insert (instruction leaflet) to your physician, pharmacist, or registered salesperson for a consultation.

Related area	Symptom
Skin	Rash/redness, itching, oedema
Gastrointestinal system	Nausea and vomiting, constipation, loss of appetite, feeling of discomfort in the stomach
Neuropsychiatric system	Dizziness, headache

3. When symptoms do not improve even after taking this medicine for 5 to 6 days, stop taking this medicine and consult a physician, dentist, pharmacist or registered salesperson, being sure to take this package insert (instruction leaflet) with you.

[Indications]

Soft stool, diarrhea, food poisoning, water poisoning, diarrhea and vomiting, loose bowels and diarrhea due to indigestion

[Dosage and Directions]

Take the following dosages with cold or warm water within 30 minutes after a meal.

Age	One dose	Daily dosage
Adults (15 years and older)	4 tablets	3 times
11-14 years of age	3 tablets	
5-10 years of age	2 tablets	
Under 5 years of age	Do not use	

[Dosage and Administration Precautions]

- (1) Follow the dosage and directions.
- (2) When allowing children 5 years or over to take the medicine, a guardian must be present, and caution must be exercised to prevent them from taking an incorrect dose or choking on it.

[Ingredients and Contents]

Every 12 tablets (maximum daily adult dosage) include the following ingredients.

Active Ingredients	Content per tablet	Effect
JP wood creosote*	270 mg	Normalization of excessive colonic motility, regulation of intestinal water content
JP powdered geranium herb	300 mg	Improvement of intestinal motility
Dried phellodendron bark essence	300 mg	Improvement of gastric function, bacteriostasis inside the intestines

[Inactive ingredients] Magnesium aluminum silicate, aluminum silicate, methyl cellulose, CMC-Ca, hydroxypropyl cellulose and magnesium stearate. The sugar coating contains white sugar, gum arabic, gelatin, talc, calcium carbonate, titanium dioxide, CMC-Na, macrogol, carnauba wax, and white beeswax.

*There are two types of substances generally called "creosote": one is JP wood-tar creosote derived from beechwood trees, and the other is industrial creosote (coal-tar creosote) extracted from coal. However, these substances are completely different in terms of raw materials, components, and use. In addition, the results of carcinogenicity tests have demonstrated that JP wood creosote is not identified as carcinogenic.

For more information or inquiries, please visit our website (<http://www.seirogan.co.jp/>) or contact our customer service department.

[Storage and Handling Precautions]

1. Keep out of reach of children.
2. Do not transfer the medicine to a different container. (It may lead to misuse or alter the quality of the drug.)
3. Store in a cool place away from direct sunlight and keep it tightly closed.
4. Do not use after the use-by date indicated on the package.

Marketing authorization holder

TAIKO PHARMACEUTICAL CO., LTD.

3-34-14, Uchihonmachi, Suita Osaka, Japan 564-0032

01004102